

RASCAL'S MEDICAL HISTORY

A5785498 49.20LBS DOG TRICOLOR N AUST CATTLE DOG

6/25/2026

INTAKE EXAM NORMAL 49.20LBS

Treatment Description:

Dog FAS Intake Medical/Behavior Form
S:
Patient was brought in as Stray, OTC
Microchip not found
O:
Behavioral Observations
Offer tasty food: No interest
Type of food offered: Canned
Overall body posture: Crouched , Reluctant to approach Frozen
Ears: Neutral
Eyes: Whale eye
Mouth: Closed
Vocalizing: Quiet
Fear-Anxiety-Stress level observed at intake: (3) Moderate FAS
Other behavior observations: nervous, scared, pillable, not leash trained- won't walk
Medical Observations:
WT 49.2 lbs
Age: ~1 year
Gender: male
BCS 5 / 9
MM- pink CRT- < 2 sec BAR
EENT: NSF
CV: NSF
RESP: NSF
Abdomen/UG: two palpable testicles
PLN: NSF
Musculoskeletal: amb x 4
Integument: NSF
Neuro: NSF
P: Routine vaccinations:
Nobivac DHLPP SQ RFL (booster in 2-4 weeks)
Bordetella Intratrach3 IN once (good for 1 year)
Bravecto 44-88lbs PO once
4.9 cc Pyrantel pamoate suspension 50mg/ml PO once
Was Entered By:

4.90 DEWORM PYRANTEL 1.00 TIMES/DAY FOR 1.00 DAYS	Date Given:6/25/2026
BRAVECTO CANINE 44-88LBS	Date Given:6/25/2026
BORDETELLA	Date Given:6/25/2026
DHLPP	Date Given:6/25/2026

7/2/2026

SX EXAM

NORMAL

45.60LBS

Treatment Description:

NOBIVAC 3 Serial number: 856741 EXP 01/28/27

Microchip applied # 956000019970455

Rx Vetprofen 100 mg tablet SID for 3 days (Pain Medication)

S: Patient presents for Dog Castration.

O: Physical examination

Temperature- 101.6 Pulse- 120 Respiration- 40 MM- pink CRT- < 2 sec BAR

Weight 45.6 lbs BCS 5 /9

EENT- mild dental tartar

Cardiovascular/Respiratory- NSF

Abdomen- NSF

Musculoskeletal- NSF

Neurologic- NSF

Lymph nodes- NSF

Genito-urinary- NSF

Integument- NSF

Behavior: very tense, nervous, whale eyed, resisted being examined, struggled to muzzle, needs BET assessment

A: Patient is fit for surgery

P: Surgery Report- Dog Neuter

Anesthesia

Pre-anesthetic: Acepromazine (2mg/ml) 0.5 cc IM, Butorphanol 10mg/ml: 0.6 cc IM,

Ampicillin 200mg/ml: 2.3 cc SQ

Induction: Ketamine (100 mg/ml) 1.0 cc IV, Midazolam (5 mg/ml) 1.0 cc IV

Intubation: Endotracheal tube size 7.5

Maintenance: Oxygen 1L Isoflurane 2%

Monitoring: Pulse oximeter

Supportive care: Thermoregulation pad/blanket

A testicle was pushed up in the scrotum on the median raphe area and a slightly off midline incision was made over the testicle (#10). The testicle was then exteriorized and freed of subcutaneous and connective tissue, manually with gauze. A carnalt was then placed over the vas deferens, cremaster, and vessels proximal to the epididymus. A single ligature (Miller's knot, 2-0 PDS) was placed proximal to the clamp and the testicle removed distal to the clamp. The clamp was then removed and the stump checked for bleeding. This procedure was repeated on the second testicle. The subcutaneous tissue was apposed using a single cruciate suture of 2-0 Monocryl. Tissue glue was placed on the skin.

Intradermal tattoo placed

Surgeon:

VETPROFEN 100

Date Given:7/2/2026

RABIES

Date Given:7/2/2026

A5785498 45.60LBS DOG TRICOLOR N AUST CATTLE DOG

IF YOU OR YOUR VETERINARIAN HAVE ANY QUESTIONS REGARDING THE MEDICAL TREATMENT YOUR
ANIMAL HAS RECEIVED
PLEASE CALL US AT (562) 940-6898