

CHARLIE'S MEDICAL HISTORY

A5762133 56.60LBS DOG BLACK / WHITE N GERM SHEPHERD MIX

3/13/2026

INTAKE EXAM

NORMAL

56.60LBS

Treatment Description:

Dog FAS Intake Medical/Behavior Form

S:

Patient was brought in as Stray

Microchip not found

O:

Behavioral Observations

Offer tasty food: No interest

Type of food offered: Canned

Overall body posture: Reluctant to approach, Stiff/tense , Frozen

Ears: Flattened back

Eyes: Avoids contact

Mouth: Closed

Vocalizing: Quiet

Fear-Anxiety-Stress level observed at intake: (3) Moderate FAS

Other behavior observations:

Medical Observations:

WT 56.6 lbs

Age: ~ 3 years

Gender: Male Intact

BCS 5 / 9

MM- pink CRT- uto BAR

EENT: Limited exam- NSF

CV: uto

RESP: uto

Abdomen/UG: uto

PLN: uto

Musculoskeletal: amb x 4

Integument: NSF

Neuro: uto

P: Routine vaccinations:

Nobivac DHLPP SQ RFL (booster in 2-4 weeks)

Bordetella Intratrach3 IN once (good for 1 year)

Bravecto 44-66lb PO once

5.7cc Pyrantel pamoate suspension 50mg/ml PO once

Person performing exam:

Assistar*

BRAVECTO CANINE 44-88LBS

Date Given:3/13/2026

5.70 DEWORM PYRANTEL 1.00
TIMES/DAY FOR 1.00 DAYS

Date Given:3/13/2026

BORDETELLA

Date Given:3/13/2026

DHLPP

Date Given:3/13/2026

3/18/2026

SX EXAM

NORMAL

56.20LBS

Treatment Description:

NOBIVAC 3 Serial number: 838919 Exp 10/31/26

Microchip applied # 956000019957109

Rx Vetprofen 100 mg tablet SID for 3 days (Pain Medication)

S: Patient presents for Dog Castration.

O: Physical examination

Temperature- 101.0 Pulse- 120 Respiration- 40 MM- pink CRT- < 2 sec BAR

Weight 56.2 lbs BCS 5/9

EENT- mod. dental calc.

Cardiovascular/Respiratory- NSF

Abdomen- NSF

Musculoskeletal- NSF

Neurologic- NSF

Lymph nodes- NSF

Genito-urinary- NSF

Integument- bi-lateral callous on elbows

Behavior: tense, appears nervous, intermittent wagging of tail, watchful, no aggression, OK to work with, scared to walk outside, muzzled out of percaution

A: Patient is fit for surgery

P: Surgery Report- Dog Neuter

Anesthesia

Pre-anesthetic: Acepromazine (2mg/ml) 0.65 cc IM, Butorphanol 10mg/ml: 0.75 cc IM,

Ampicillin 200mg/ml: 2.81 cc SQ

Induction: Ketamine (100 mg/ml) 1.2 cc IV, Midazolam (5 mg/ml) 1.2 cc IV

Intubation: Endotracheal tube size 10

Maintenance: Oxygen 1L Isoflurane 2%

Monitoring: Pulse oximeter

Supportive care: Thermoregulation pad/blanket

A testicle was pushed up in the scrotum on the median raphe area and a slightly off midline incision was made over the testicle (#10). The testicle was then exteriorized and freed of subcutaneous and connective tissue, manually with gauze. A carnalt was then placed over the vas deferens, cremaster, and vessels proximal to the epididymus. A single ligature (Miller's knot, 2-0 PDS) was placed proximal to the clamp and the testicle removed distal to the clamp. The clamp was then removed and the stump checked for bleeding. This procedure was repeated on the second testicle. The subcutaneous tissue was apposed using a single cruciate suture of 2-0 Monocryl. Tissue glue was placed on the skin.

Intradermal tattoo placed

Surgeon:

RVT:

VETPROFEN 100

Date Given:3/18/2026

RABIES

Date Given:3/18/2026

3/18/2026

RVT EXAM NORMAL

Treatment Description:

Emailed BET regarding patients behavior and possible behavior assessmnet.

SX EXAM NORMAL 56.20LBS

Treatment Description:

Sx Addendum: Alteration Sx done by FY 495 3/18/26

IF YOU OR YOUR VETERINARIAN HAVE ANY QUESTIONS REGARDING THE MEDICAL TREATMENT YOUR
ANIMAL HAS RECEIVED
PLEASE CALL US AT (562) 940-6898