

HAL'S MEDICAL HISTORY

A5762959 75.60LBS DOG BLACK / TAN N GERM SHEPHERD

3/17/2026

INTAKE EXAM

OTHER

75.60LBS

Treatment Description:

Dog FAS Intake Medical/Behavior Form

S:

Patient was brought in as Stray, OTC

Microchip not found

O:

Behavioral Observations

Offer tasty food: No interest

Type of food offered: Canned

Overall body posture: Upright , Stiff/tense

Ears: Neutral

Eyes: Whale eye

Mouth: Closed

Vocalizing: Quiet

Fear-Anxiety-Stress level observed at intake: (3) Moderate FAS

Other behavior observations: nervous, scared

Medical Observations:

WT 75.6 lbs

Age: ~6 years

Gender: female

BCS 6 / 9

MM- pink CRT- < 2 sec BAR

EENT: fly strikes AU

CV: NSF

RESP: NSF

Abdomen/UG: NSF

PLN: NSF

Musculoskeletal: amb x 4

Integument: scars on face, slight swelling/abrasion to right side of muzzle

Neuro: NSF

P: Routine vaccinations:

Nobivac DHLPP SQ RFL (booster in 2-4 weeks)

Bordetella Intratrach3 IN once (good for 1 year)

bravecto 44-88lbs PO once

7.6 cc Pyrantel pamoate suspension 50mg/ml PO once

Cephalexin 500mg: 1.5 cap po bid x 7d

Vetprofen 100mg: 1.5 tab po sid x 5d

Vet to recheck

Was Entered By: -----

Exam/Treatment performed by :

Assisted by:

1.50 VETPROFEN 100 1.00
TIMES/DAY FOR 5.00 DAYS

Date Given:3/17/2026

1.50 CEPHALEXI 500 2.00
TIMES/DAY FOR 7.00 DAYS

Date Given:3/17/2026

7.60 DEWORM PYRANTEL 1.00
TIMES/DAY FOR 1.00 DAYS

Date Given:3/17/2026

3/17/2026

BRAVECTO CANINE 44-88LBS Date Given:3/17/2026
BORDETELLA Date Given:3/17/2026
DHLPP Date Given:3/17/2026

3/18/2026

VET EXAM OTHER

Treatment Description:

NIC

3/19/2026

VET EXAM OTHER

Treatment Description:

S: On vet check for check skin
O: PHYSICAL EXAM 10 mins
HR: 130 Respiration: pant CRT <2s BAR MM Pink BCS 4/9
Patient:
EENT: eyes clear OU, nose clean, ears grossly clean AU, mild dental calc
LN: nsf
Thorax: eupneic, no obvious signs of distress, no apparent
Abdomen: n/e
Genitourinary: no obvious lesions/discharge noted
Integument: mild ear tip crusting AU, no swelling of the muzzle
Musculoskeletal: amb x4
Neuro: appropriate mentation, no gross deficits

A: Fly strike lesions
R muzzle swelling appears to be resolved

P: Ok to cont txt per SP
Monitor for appetite, behavior and worsening condition and contact veterinarian if this occurs. Feed canned food to entice appetite, if necessary.
HEALTH WAIVER

3/24/2026

3/24/2026

SX EXAM

NORMAL

67.00LBS

Treatment Description:

NOBIVAC 3 Serial number: 838919 EXP 10/31/26

Microchip applied # 956000019932273

Rx Vetprofen 100 mg 1.5 tablet SID for 3 days (Pain Medication)

S: Patient presents for Dog Castration/Right Abdominal Cryptorchid.

O: Physical examination

Temperature- 100.8 Pulse- 120 Respiration- pant MM- pink CRT- < 2 sec BAR

Weight 67 lbs BCS 5/9

EENT- mod. dental calc.

Cardiovascular/Respiratory- NSF

Abdomen- NSF

Musculoskeletal- NSF

Neurologic- NSF

Lymph nodes- NSF

Genito-urinary- abdominally retained right testicle

Integument- NSF

Behavior: friendly, moves around a lot, no aggression, Ok to work with

A: Patient is apparently healthy.

P: Surgery Report- Dog Neuter

Anesthesia

Pre-anesthetic: Acepromazine (2mg/ml) 0.55 cc IM, Hydromorphone 2mg/ml: 0.9 cc IM,

Ampicillin 200mg/ml 3.35 cc SQ

Induction: Ketamine (100 mg/ml) 1.4 cc IV, Midazolam (5 mg/ml) 1.4 cc IV

Intubation: Endotracheal tube size 10

Maintenance: Oxygen 2L Isoflurane 2%

Monitoring: Pulse oximeter

Supportive care: Thermoregulation pad/blanket

A 2 cm paramedian incision was made just lateral to the cranial prepuce with a 10 blade. The subcutaneous tissues with bluntly dissected with metzenbaum scissors. The sheath of the rectus was identified and incised with a 10 blade. The external and internal rectus muscles were gently separated using carnalms until the peritoneum was visualized. The peritoneum was tented and a stab incision was made to enter the abdomen. The abdominal testicle was identified and a carnalm was then placed over the vas deferens, cremaster, and vessels proximal to the epididymus. A single ligature (Miller's knot, 2-0 PDS) was placed proximal to the clamp and the testicle removed distal to the clamp. The clamp was then removed and the stump checked for bleeding. The rectus sheath was closed using 2-0 monocryl in a simple continuous pattern. The subcutaneous and intradermal layers were closed using 2-0 monocryl in a simple continuous pattern. Tissue glue was placed on the skin. The remaining testicle was removed in a routine fashion.

Interdermal tattoo placed

Surgeo

1.50 VETPROFEN 100 1.00
TIMES/DAY FOR 3.00 DAYS

Date Given:3/24/2026

RABIES

Date Given:3/24/2026

3/24/2026

SX EXAM

NORMAL

53.40LBS

Treatment Description:

NOBIVAC 3 Serial number: 838919 Exp 10/31/26

Microchip applied # 956000019931908

Rx Vetprofen 100 mg tablet SID for 3 days (Pain Medication)

S: Patient presents for Dog Castration.

O: Physical examination

Temperature- 100.5 Pulse- 120 Respiration- pant MM- pink CRT- < 2 sec BAR

Weight 53.4lbs BCS 5/9

EENT- NSF

Cardiovascular/Respiratory- NSF

Abdomen- NSF

Musculoskeletal- NSF

Neurologic- NSF

Lymph nodes- NSF

Genito-urinary- NSF

Integument- NSF

Behavior: friendly, easy to work with

A: Patient is fit for surgery

P: Surgery Report- Dog Neuter

Anesthesia

Pre-anesthetic: Acepromazine (2mg/ml) 0.6 cc IM, Butorphanol 10mg/ml: 0.7 cc IM,

Ampicillin 200mg/ml: 2.7 cc SQ

Induction: Ketamine (100 mg/ml) 1.2 cc IV, Midazolam (5 mg/ml) 1.2 cc IV

Intubation: Endotracheal tube size 10

Maintenance: Oxygen 1L Isoflurane 2%

Monitoring: Pulse oximeter

Supportive care: Thermoregulation pad/blanket

A testicle was pushed up in the scrotum on the median raphe area and a slightly off midline incision was made over the testicle (#10). The testicle was then exteriorized and freed of subcutaneous and connective tissue, manually with gauze. A Carmalt was then placed over the vas deferens, cremaster, and vessels proximal to the epididymus. A single ligature (Miller's knot, 2-0 PDS) was placed proximal to the clamp and the testicle removed distal to the clamp. The clamp was then removed and the stump checked for bleeding. This procedure was repeated on the second testicle. The subcutaneous tissue was apposed using a single cruciate suture of 2-0 Monocryl. Tissue glue was placed on the skin.

Intradermal tattoo placed

Cerenia inj 10 mg/ml 2.4 ml SQ- VO at time of extubation

Surgeon: RVT:

VETPROFEN 100

Date Given:3/24/2026

RABIES

Date Given:3/24/2026

3/24/2026

RVT EXAM

NORMAL

Treatment Description:

Omit- T26-557276

Incorrect Entry

RVT:

IF YOU OR YOUR VETERINARIAN HAVE ANY QUESTIONS REGARDING THE MEDICAL TREATMENT YOUR
ANIMAL HAS RECEIVED
PLEASE CALL US AT (562) 940-6898